

208517

3 05736 55204 6



READ AND KEEP CARTON FOR COMPLETE  
WARNINGS AND INFORMATION

### Drug Facts

**Active ingredients (in each capsule)** **Purposes**  
Diphenhydramine hydrochloride 25 mg..... Nighttime sleep-aid  
Solubilized ibuprofen equal to  
200 mg ibuprofen (NSAID)\* ..... Pain reliever  
(present as the free acid and potassium salt)  
\*nonsteroidal anti-inflammatory drug

**Uses** ■ for relief of occasional sleeplessness when  
associated with minor aches and pains ■ helps you fall  
asleep and stay asleep

### Warnings

**Allergy alert:** Ibuprofen may cause a severe allergic reaction,  
especially in people allergic to aspirin. Symptoms may include:  
■ hives ■ facial swelling ■ asthma (wheezing) ■ shock  
■ skin reddening ■ rash ■ blisters  
If an allergic reaction occurs, stop use and seek medical  
help right away.

**Stomach bleeding warning:** This product contains an  
NSAID, which may cause severe stomach bleeding. The  
chance is higher if you ■ are age 60 or older ■ have had  
stomach ulcers or bleeding problems ■ take a blood  
thinning (anticoagulant) or steroid drug ■ take other drugs  
containing prescription or nonprescription NSAIDs [aspirin,  
ibuprofen, naproxen, or others] ■ have 3 or more alcoholic  
drinks every day while using this product ■ take more or  
for a longer time than directed

### Drug Facts (continued)

**Heart attack and stroke warning:** NSAIDs, except aspirin,  
increase the risk of heart attack, heart failure, and stroke.  
These can be fatal. The risk is higher if you use more than  
directed or for longer than directed.

**Do not use** ■ if you have ever had an allergic reaction to  
any other pain reliever/fever reducer ■ unless you have  
time for a full night's sleep ■ in children under 12 years of  
age ■ right before or after heart surgery ■ with any other  
product containing diphenhydramine, even one used on skin  
■ if you have sleeplessness without pain

**Ask a doctor before use if** ■ stomach bleeding warning  
applies to you ■ you have problems or serious side effects  
from taking pain relievers or fever reducers ■ you have a  
history of stomach problems, such as heartburn ■ you  
have high blood pressure, heart disease, liver cirrhosis,  
kidney disease, asthma, or had a stroke ■ you are taking a  
diuretic ■ you have a breathing problem such as emphysema  
or chronic bronchitis ■ you have glaucoma ■ you have  
trouble urinating due to an enlarged prostate gland

**Ask a doctor or pharmacist before use if you are**  
■ taking sedatives or tranquilizers, or any other sleep-aid  
■ under a doctor's care for any continuing medical illness  
■ taking any other antihistamines ■ taking aspirin for  
heart attack or stroke, because ibuprofen may decrease this  
benefit of aspirin ■ taking any other drug

**When using this product** ■ drowsiness will occur  
■ avoid alcoholic drinks ■ do not drive a motor vehicle or  
operate machinery ■ take with food or milk if stomach  
upset occurs

### Drug Facts (continued)

#### Stop use and ask a doctor if

■ you experience any of the following signs of  
stomach bleeding:  
■ feel faint ■ vomit blood ■ have bloody or black  
stools ■ have stomach pain that does not get better  
■ you have symptoms of heart problems or stroke:  
■ chest pain ■ trouble breathing ■ weakness in one  
part or side of body ■ slurred speech ■ leg swelling  
■ pain gets worse or lasts more than 10 days  
■ sleeplessness persists continuously for more than  
2 weeks. Insomnia may be a symptom of a serious underlying  
medical illness. ■ redness or swelling is present in the  
painful area ■ any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional  
before use. It is especially important not to use ibuprofen at  
20 weeks or later in pregnancy unless definitely directed to do  
so by a doctor because it may cause problems in the unborn  
child or complications during delivery. **Keep out of reach of  
children.** In case of overdose, get medical help or contact a  
Poison Control Center right away.

**Directions** ■ do not take more than directed  
■ adults and children 12 years and over: take 2 capsules  
at bedtime ■ do not take more than 2 capsules in 24 hours

#### Other information

■ each capsule contains: **potassium 20 mg**  
■ read all warnings and directions before use. Keep carton.  
■ store at 20-25°C (68-77°F)

### Drug Facts (continued)

■ avoid excessive heat above 40°C (104°F)  
■ protect from light

**Inactive ingredients:** D&C red no. 33, FD&C blue no. 1,  
gelatin, medium-chain triglycerides, pharmaceutical ink,  
polyethylene glycol, potassium hydroxide, purified water,  
sorbitol sorbitan solution

#### Questions or comments?

Call weekdays 9 AM to 5 PM EST at 1-800-88-ADWIL

**Do Not Use if seal under bottle cap imprinted with  
"SEALED FOR YOUR PROTECTION" is broken or missing.**

For most recent product information, visit [www.adwll.com](http://www.adwll.com)  
Trademarks are owned by Haleson, Warren, NJ 07059.  
©2012 Haleson group of companies or its licensor.  
L001-49E-516 at Haleson or registered trademark  
of Haleson Pharma Solutions.

128 CODE  
AREA128 CODE  
AREA

HALESON

NEW  
Smaller  
Capsule  
Same Strength\*

# Advil PM<sup>®</sup>

## Liqui-Gels<sup>®</sup>

Solubilized ibuprofen, 200 mg/  
Diphenhydramine HCl, 25 mg  
Pain Reliever (NSAID)/  
Nighttime Sleep-Aid



40  
Liquid Filled  
Capsules

Actual  
Size

\*Compared to Advil PM Liqui-Gels

128 CODE  
AREA128 CODE  
AREA